

## Recenze článků do odborných časopisů – Jan Riegert

**2021**

Yetis, R., Nergiz, H. & Yetis, A. D. Effects of water quality on the species richness and population distribution of waterbirds in Ahlat Marshes, Turkey (**Biologia**)

Klaus, S. & Ludwig, T. Long-term trends of Hazel Grouse (*Tetrastes bonasia*) in the Bohemian Forest (Šumava), Czech Republic, 1972-2019 (**Birds**)

[Double-blind system] Long term changes in a small mammal community under multiple disturbances (**Integrative Zoology**)

Iezekiel, S., Yosef, R., Themistokleus, C., Bakaloudis, D. E., Vlachos, Ch. G., Antoniou, A., Iezekiel, E., Papakosta, M. A. & Kosicki, J. Z. Endemic Cyprus Scops Owl *Otus cyprius* readily adapts to artificial nest boxes (**Animals**)

Tavankar, F., Latterini, F., Nikooy, M., Venanzi, R., Naghdi, R., Picchio, R. Forest management and silvicultural treatments influence on abundance of snags and bird cavity-nests in mountain mixed beech forests (**Environments**)

Amo, L. & Saavedra, I. Attraction to smelly food in birds: Insectivorous birds discriminate between the pheromones of their prey and those of non-prey insects (**Biology**)

Abeyta, E. J., Bartlow, A. W., Hathcock, Ch. D. & Fair, J. M. Individual nest site preferences do not explain upslope population shifts of a secondary cavity-nesting species (**Animals**)

[Double-blind system] Distance to range edge and forest cover determine intraspecific variation in sensitivity to habitat fragmentation (**Global Ecology and Biogeography**)

Binda1, V. A., Kenfack, I. B., Dong, A. B. E., Gandiwa, E. & Mayaka, T. B. Enchantment on Mount Oku (Cameroon) Farm Landscape: Is there a gradient in avian diversity metrics? (**Frontiers in Ecology and Evolution, section Biogeography and Macroecology**)

[Double-blind system] Large modern oases are key sites for Eurasian scops owl in Tunisian pre-Saharan: implications for wildlife management in dry farmlands (**Biologia**)

Kaplan, G. Casting the Net Widely for Change in Animal Welfare: The Plight of Birds in Zoos, Ex-Situ Conservation and Conservation Fieldwork (**Animals**)

[Double-blind system] Reed die-back and conservation of small reed birds at Lake Neusiedl, Austria (**Biologia**)

Bruna, P. & Monticelli, P. Maned wolf duet? The first record of two maned wolves vocalizing simultaneous (**Academia Letters**)

Liu, J. & Liang, W. Hiss matters: Hissing calls of tits elicit vigilance in feeding squirrels (**Ethology Ecology & Evolution**)

Mettke-Hofmann, C. Morph composition matters in the Gouldian finch: Involvement of red-headed birds increases vigilance (**Birds**)

Albayra, T. & Ahmet İhsan Aytek, A. I. Bill variation of captive and wild Chukar Partridge populations: shape or size (**Diversity**)

## **2020**

Sliwinski, K., Ronnenberg, K., Jung, K., Strauß, E. & Siebert, U. Habitat requirements of the European brown hare (*Lepus europeus* Pallas 1778) in an intensively used agriculture region (Lower Saxony, Germany) (**BMC Ecology**)

Sakhvon, V. & Kövér, L. Distribution and habitat preferences of the urban Woodpigeon (*Columba palumbus*) in the north-eastern breeding range in Belarus (**Landscape and Urban Planning**)

Johann, F. & Arnold, J. Scattered woody vegetation promotes European brown hare (*Lepus europaeus*) population (**Basic and Applied Ecology**)

Szymański, P., Olszowiak, K., Wheeldon, A., Budka, M. & Osiejuk, T. S. Passive acoustic monitoring gives new insight into year-round duetting behaviour of a tropical songbird (**Ecological Indicators**)

Wheeldon, A., Szymański, P. & Osiejuk, T. S. Yellow-breasted Boubous (*Laniarius atroflavus*) jointly defend territories with male led duets against stranger pairs, males and females (**Ethology**)

Shivambu, C. T., Shivambu, N., Nelufule, T., Moshobane, M. C., Zungu, M. M. & Downs, C. T. Numbers of invasive House Sparrows (*Passer domesticus* Linnaeus, 1758) in a rural landscape of Limpopo Province, South Africa (**Ostrich**)

Yetis, R., Nergiz, H. & Yetis, A. D. Effects of water quality on the species richness and population distribution of waterbirds in Ahlat Marshes, Turkey (**Biologia**)

Prawesthi, G. A. & Hadisusantho, S. Ecological niche differentiation of three species of Nectariniidae in agroforestry areas of Jatimulyo village, Yogyakarta, Indonesia (**KUKILA**)

Šotnár, K., Obuch, J., Pačenovský, S & Jarčuška, B. Spatial distribution of four sympatric owl species in a Carpathian montane forests, Slovakia (**Slovak Raptor Journal**)

Kubelka, V., Havlíček, J. & Mikeš, V. Zajímavá a vzácná ornitologická pozorování v jižních Čechách VII (**Sborník Jihočeského Muzea v Českých Budějovicích, Přírodní vědy**)

**2019**

Sliwinski, K., Ronnenberg, K., Jung, K., Strauß, E. & Siebert, U. Habitat requirements of the European brown hare (*Lepus europeus* Pallas 1778) in an intensively used agriculture region (Lower Saxony, Germany) (**BMC Ecology**)

Sakhvon, V. & Kövér, L. Distribution and habitat preferences of the urban Woodpigeon (*Columba palumbus*) in the north-eastern breeding range in Belarus (**Landscape and Urban Planning**)

Johann, F. & Arnold, J. Scattered woody vegetation promotes European brown hare (*Lepus europaeus*) population (**Basic and Applied Ecology**)

Szymański, P., Olszowiak, K., Wheeldon, A., Budka, M. & Osiejuk, T. S. Passive acoustic monitoring gives new insight into year-round duetting behaviour of a tropical songbird (**Ecological Indicators**)

Wheeldon, A., Szymański, P. & Osiejuk, T. S. Yellow-breasted Boubous (*Laniarius atroflavus*) jointly defend territories with male led duets against stranger pairs, males and females (**Ethology**)

Shivambu, C. T., Shivambu, N., Nelufule, T., Moshobane, M. C., Zungu, M. M. & Downs, C. T. Numbers of invasive House Sparrows (*Passer domesticus* Linnaeus, 1758) in a rural landscape of Limpopo Province, South Africa (**Ostrich**)

Yetis, R., Nergiz, H. & Yetis, A. D. Effects of water quality on the species richness and population distribution of waterbirds in Ahlat Marshes, Turkey (**Biologia**)

Zajímavá a vzácná ornitologická pozorování v jižních Čechách VI (**Sborník Jihočeského Muzea v Českých Budějovicích**)

**2018**

[Double-blind review] Benefits of extra food to reproduction depend on maternal condition (**Oikos**)

Gryz, J. & Gryz, D. K. Changes in the tawny owl *Strix aluco* diet along an urbanisation gradient  
**(Biologia)**

Weller, A. A. First record of subsong and vocal duetting in the Purple Sunbird (*Cinnyris asiaticus*)  
**(Sandgrouse)**

Závora, J. & Pykal, J. Pravidelné přezimování sokola stěhovavého tundrového (*Falco peregrinus calidus*) v Českých Budějovicích v letech 2013-2017 (**Sborník Jihočeského muzea v Českých Budějovicích**)

## **2017**

[Double-blind system] Roosting behaviour is related to reproductive strategy in brood parasitic cowbirds (**Ibis**)

[Double-blind system] A high incidence of non-cavity nesting in an introduced population of House Sparrows suggests that the species should not be constrained by cavity-nest site availability  
**(Avian Research)**

[Double-blind system] The effect of industrial noise on owl occupancy and seasonal use in the boreal forest (**Avian Conservation and Ecology**)

## **2016**

Kreiderits, A., Gamauf, A., Krenn H. W., Sumasgutner, P. The influence of local weather conditions and alternative prey composition on the breeding performance of urban Eurasian kestrels (*Falco tinnunculus*) (**Bird Study**)

Veselovský, T., Bacsa, K. & Tulis, F. Barn Owl (*Tyto alba*) diet composition on intensively used agricultural land in the Danube Lowland (**Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis**)

Nguembock, B., Sali, M., William, K., Olivia, A. E. D., Yollande, G., Clinton, F. & Herve, N. D. P. Bird survey of the Abobo-Etetak hill (Yaounde, Cameroon) and a glance on the avifaunistic diversity of this hill (Abobo-Etetak) (**The Open Ornithology Journal**)